California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Race/Ethnicity and Year		E	Birthweight	(4)			Hei	ght and V	Veight			Aner	nio
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Low H	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th	Number	%
White, Not Hispanic	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	16,347 15,825 13,214 17,421 19,395 21,696 23,690 28,173 30,187 30,249	7.8 7.0 6.8 7.6 8.4 7.9 8.3 8.6 10.9 11.5	10.2 10.4 10.7 11.0 11.1 11.1 10.3 10.6 10.4 10.2	77,990 79,702 74,176 90,070 95,308 109,489 119,187 125,312 130,944 135,322	4.8 4.8 5.1 5.0 5.1 4.9 5.1 5.1 5.3	7.2 7.0 7.5 7.3 7.4 7.3 7.7 7.7 7.2 7.3	12.3 12.3 11.9 12.1 11.7 11.4 10.8 10.7 11.1 10.7	27,919 28,704 26,702 34,273 36,095 42,390 45,172 44,894 45,887 48,830	15.3 15.2 15.5 14.9 14.8 15.3 14.5 14.4	13.1 13.4 12.3 12.8 12.0 11.5 10.7 10.7 10.5 10.2	37,832 38,533 35,973 46,577 51,674 60,197 64,147 72,208 75,868 79,778	12.2 11.7 12.6 11.8 11.6 12.1 12.3 13.1 13.4 13.1
Black, Not Hispanic	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	8,401 8,217 6,704 9,584 10,304 11,241 12,662 13,364 14,449 15,083	12.1 12.2 11.6 11.6 11.7 12.0 11.7 12.6 13.4 14.9	5.4 6.3 6.3 5.9 5.9 5.9 5.9 5.9 5.7	47,529 47,190 44,539 56,062 59,230 66,764 75,850 75,471 80,413 79,846	6.4 6.6 6.8 7.0 7.0 6.8 7.0 7.0 7.1	6.6 6.4 6.6 6.4 6.5 6.6 6.9 7.0 6.9 7.1	14.3 14.3 14.2 14.4 13.8 13.3 12.6 12.3 12.4 12.4	19,609 19,215 18,073 23,531 25,036 28,511 32,966 31,794 32,625 31,716	14.4 14.2 14.3 14.1 14.6 14.1 13.2 13.1 12.8 12.8	12.9 13.0 12.2 12.5 11.5 11.1 10.2 10.2 10.1 9.6	23,882 22,962 21,801 29,602 32,659 36,849 42,148 44,698 46,072 46,004	17.7 16.5 18.3 17.8 17.7 18.1 18.5 19.7 20.8 20.0



Page:

Run Date:

5/5/04

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Surveillance (1) Run Date:

Page:

5/5/04

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity Children Aged < 5 Years (2)

	Sumi	mary of T	rends ir	n Growt	h and An	emia In	dicato	rs by F	Race/Eth	nicity (3)			
Race/Ethnicity and Year		В	irthweight	(4)		Height and Weight							nia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Low Hk (7)	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% n <u>≥</u> 95th	Number	%
Hispanic	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	115,362 112,084 93,404 114,019 112,491 114,612 123,572 135,656 134,728 139,685	6.6 6.1 6.1 6.2 6.1 6.0 6.1 6.2 7.8 8.4	8.7 9.3 9.4 9.7 9.9 9.7 9.6 9.7 9.5 9.4	588,054 583,889 546,974 616,504 616,024 648,721 678,216 664,345 651,808 633,719	4.7 4.7 4.8 4.7 4.5 4.5 4.6 4.6	4.7 4.7 4.6 4.5 4.5 4.8 4.8 4.7	17.7 17.5 17.0 17.3 16.9 16.7 16.0 15.8 16.0	242,274 237,783 221,484 258,491 264,862 275,060 285,600 272,638 255,373 237,513	16.7 16.9 16.7 16.7 16.8 16.9 16.2 16.3 16.2 16.3	19.0 18.6 17.7 17.8 17.1 16.8 16.1 15.9 16.0 15.9	374,196 363,388 340,440 412,834 429,996 450,480 469,530 483,187 461,491 443,054	13.6 13.0 13.4 13.1 12.6 12.6 13.1 13.4 14.0 14.2
American Indian/Alaskan Native	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	381 362 384 484 603 532 535 623 708 703	8.4 6.1 4.7 5.6 6.5 3.9 7.1 5.9 9.0	13.6 11.0 8.9 10.3 10.8 15.8 11.2 10.3 11.3	2,231 2,340 2,521 3,062 3,413 3,147 3,222 3,402 3,265 3,351	6.3 5.0 6.9 4.9 5.7 4.7 5.2 4.6 5.2	5.8 5.6 5.1 6.0 5.0 5.3 5.1 5.9 5.5	18.1 19.3 18.6 17.0 16.4 16.7 17.0 16.3 16.7 16.4	872 959 1,007 1,282 1,387 1,370 1,403 1,403 1,301 1,307	15.7 19.5 20.2 16.4 17.9 19.6 18.5 18.0 17.0	21.8 20.8 19.7 17.5 17.5 18.2 17.7 16.4 19.6 16.0	1,324 1,341 1,489 1,963 2,208 2,101 2,071 2,381 2,485 2,380	11.9 12.5 15.2 13.9 15.4 17.1 16.5 14.4 14.0 13.2



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Children Aged < 5 Years (2)

California
Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity

Page: 3 Run Date: 5 / 5 / 04

Race/Ethnicity and Year		E	Birthweight	(4)				A	!-				
·			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Aner Low H (7	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% n <u>></u> 95th	Number	%
Asian/Pacific Islander	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	8,736 6,637 5,920 5,373 5,362 5,854 6,937 7,956 8,605 9,873	6.7 7.4 7.7 8.2 7.4 7.3 8.4 8.0 7.9 8.6	6.5 5.4 5.5 5.5 5.5 5.7 5.5 5.5 5.4 5.1	54,114 43,633 34,663 35,708 36,906 42,543 48,585 51,261 54,160 55,133	4.9 5.6 6.1 6.3 6.7 6.8 6.8 6.7 6.9 7.2	6.1 5.9 6.3 6.4 5.9 5.3 5.9 5.9 6.0	13.4 13.6 12.8 13.6 13.3 13.6 12.9 12.7 12.2	18,282 14,676 11,996 15,210 16,320 19,043 21,493 22,135 22,232 22,120	14.3 14.5 14.5 14.5 14.5 15.1 14.3 14.5 13.9 14.1	15.2 15.0 14.7 15.2 14.2 14.6 13.9 13.8 12.5 12.8	22,761 19,224 13,620 17,204 19,330 23,685 27,725 31,449 33,289 34,468	14.3 13.5 13.9 14.4 13.9 14.2 14.6 14.6 15.2
All Other/Unknown	2003 2002 2001 2000 1999 1998 1997 1996 1995	11,270 9,199 7,606 8,234 7,758 7,522 7,511 8,007 8,209 8,106	7.8 7.9 7.5 9.0 8.0 7.5 7.6 7.7 9.3 10.3	8.2 8.5 8.4 8.7 9.1 9.1 8.5 8.7 7.9	62,193 53,579 44,300 50,116 48,550 46,390 46,739 42,352 40,656 39,470	5.5 5.4 5.6 5.9 5.6 5.4 5.2 5.1 5.2 5.2	7.5 7.5 7.2 7.0 7.1 7.1 7.5 7.6 7.7 8.2	12.9 13.3 13.6 13.5 12.5 12.4 11.6 11.5 11.5	20,686 17,755 14,512 16,882 17,304 16,080 15,948 14,300 14,110 13,712	15.0 14.9 15.3 15.2 15.0 14.8 14.0 13.6 13.5	15.0 15.3 14.8 15.0 13.5 13.2 12.8 12.8 11.6	25,425 22,270 18,771 22,968 24,913 22,976 21,753 21,519 20,694 21,104	14.0 14.1 15.7 14.1 14.6 14.4 14.1 15.6 16.0 15.4



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

ifornia

Page:

Run Date:

5/5/04

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Race/Ethnicity and Year		Birthweight (4)				Height and Weight							nia
		Low		High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Low Hb	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th	Number	%
Total All Race/Ethnic Groups	2003 2002 2001 2000 1999 1998 1997 1996 1995 1994	160,497 152,324 127,232 155,115 155,913 161,457 174,907 193,779 196,886 203,699	7.1 6.7 6.6 6.9 6.9 6.8 7.0 7.1 8.8 9.5	8.6 9.0 9.1 9.4 9.6 9.4 9.2 9.4 9.2	832,111 810,333 747,173 851,522 859,431 917,054 971,799 962,143 961,246 946,841	4.9 4.9 5.0 5.1 5.0 4.9 4.9 5.0 5.1	5.3 5.3 5.2 5.2 5.1 5.5 5.5 5.4 5.5	16.4 16.3 16.0 16.2 15.7 15.4 14.7 14.5 14.6	329,642 319,092 293,774 349,669 361,004 382,454 402,582 387,164 371,528 355,198	16.2 16.3 16.2 16.3 16.4 15.6 15.7 15.5	17.7 17.4 16.6 16.7 15.9 15.5 14.8 14.6 14.4	485,420 467,718 432,094 531,148 560,780 596,288 627,374 655,442 639,899 626,788	13. 13. 13. 13. 13. 13. 14. 14.



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 5/5/04

Summary of Race/Ethnicity and Year	Trends in (Growth ar		nia Ind	'	y Race	Ethnicity	(3)
riaco, Emilioty and Total			Short Stature (4)	Under- weight (5)			Anem Low Hb	
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	(6) Number	%
White, Not Hispanic	2003	44,445	5.7	3.1	17.3	18.2	33,380	12.8
•	2002	46,664	6.0	3.1	16.7	17.7	34,688	12.4
	2001	42,827	5.9	3.2	16.7	16.8	31,892	12.3
	2000	57,387	6.0	3.2	16.2	16.4	40,809	11.7
	1999	61,869	5.9	3.0	16.0	15.4	45,794	10.8
	1998	62,261	5.6	3.0	16.2	14.1	46,901	10.8
	1997	60,022	5.8	3.4	15.1	13.1	45,163	11.1
	1996	51,988	5.8	3.5	15.0	12.7	44,214	12.0
	1995	50,239	6.3	3.6	14.7	12.1	44,618	12.8
	1994	49,568	6.1	3.6	14.5	11.5	44,683	12.6
Black, Not Hispanic	2003	40,349	4.3	2.7	16.8	19.3	30,544	22.3
	2002	38,428	4.1	2.6	17.2	18.8	28,859	21.0
	2001	35,294	4.0	2.8	16.5	17.5	26,428	21.1
	2000	47,248	4.1	2.9	16.4	16.6	33,569	21.5
	1999	49,495	4.2	3.1	16.1	16.2	35,394	20.1
	1998	47,538	4.1	3.4	15.4	14.8	35,232	19.9
	1997	47,805	4.2	3.9	15.1	13.8	35,527	21.0
	1996	41,375	4.2	3.8	14.8	13.1	35,591	21.0
	1995	39,418	4.5	3.8	14.4	13.1	33,127	22.3
	1994	37,059	4.4	3.8	14.2	13.0	31,372	22.1



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 5/5/04

Summary of Tren	nds in C	Growth an	d Anen	nia Ind	icators by	y Race	/Ethnicity	(3)
Race/Ethnicity and Year			Heigh	nt and W	eight			
			Short Stature (4)	Under- weight (5)			Anem Low Hb	
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	(6) Number	%
Hispanic	2003	337,291	6.1	2.3	19.4	23.2	331,214	11.5
•	2002	342,337	6.5	2.4	18.8	22.2	326,786	10.8
	2001	326,131	6.7	2.4	18.5	21.2	304,949	11.4
	2000	384,954	6.8	2.4	18.7	21.3	355,925	11.7
	1999	400,540	6.6	2.4	18.5	20.6	371,107	10.9
	1998	355,388	6.6	2.3	18.3	19.4	331,814	11.1
	1997	325,912	6.8	2.6	17.7	18.2	307,259	12.0
	1996	277,803	7.4	2.6	17.5	17.5	281,848	11.9
	1995	256,470	8.1	2.7	17.5	16.9	256,833	12.4
	1994	227,738	8.7	2.8	17.2	16.2	231,831	12.7
American Indian/Alaskan Native	2003	1,340	6.9	2.5	18.3	27.0	1,141	10.4
	2002	1,455	4.9	2.1	20.5	24.8	1,239	11.3
	2001	1,406	6.0	2.3	18.7	24.7	1,154	13.0
	2000	1,968	5.5	1.8	20.5	25.9	1,622	12.7
	1999	2,212	6.3	2.3	19.3	21.9	1,725	12.2
	1998	1,917	5.9	2.9	19.5	22.2	1,554	10.4
	1997	1,805	4.9	2.4	19.4	19.4	1,391	12.7
	1996	1,643	5.6	3.0	18.8	17.2	1,510	11.3
	1995	1,378	4.1	3.0	17.4	18.9	1,487	11.8
	1994	1,240	4.6	2.3	18.6	16.3	1,367	12.3



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 5/5/04

Summary of	Trends in (Growth ar		nia Ind		y Race	/Ethnicity	(3)
Race/Ethnicity and Year			Short Stature (4)	Under- weight (5)	Over		Anem Low Hb	
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	(6) Number	%
Asian/Pacific Islander	2003	36,958	11.4	4.8	14.0	15.2	20,517	14.4
	2002	32,276	12.3	5.0	13.8	14.2	19,110	14.0
	2001	27,026	12.6	5.1	13.7	13.2	16,326	15.1
	2000	37,206	13.7	5.3	13.0	12.9	21,827	13.3
	1999	38,569	13.2	5.4	12.6	11.6	23,711	14.1
	1998	36,892	14.5	5.7	12.1	10.6	24,786	14.7
	1997	33,631	14.6	6.3	11.1	9.9	24,067	14.6
	1996	29,441	15.7	6.5	10.5	8.9	23,704	16.2
	1995	26,394	17.7	7.4	9.9	7.8	22,423	15.9
	1994	25,189	18.9	7.6	9.5	7.3	22,660	15.1
All Other/Unknown	2003	29,631	6.8	3.4	16.9	20.8	20,826	15.2
	2002	26,326	7.3	3.5	17.4	19.3	19,020	14.6
	2001	21,665	7.2	3.4	16.3	18.3	16,506	15.4
	2000	26,448	7.1	3.7	16.5	18.3	19,541	15.1
	1999	27,090	7.5	3.8	16.3	17.1	20,665	14.7
	1998	23,061	7.8	3.9	15.5	15.2	17,878	14.5
	1997	20,764	7.8	4.9	14.8	14.1	15,800	14.2
	1996	16,746	8.4	4.7	14.7	13.1	14,577	15.9
	1995	15,238	8.5	5.3	14.6	12.4	13,237	15.7
	1994	14,352	9.1	5.4	13.7	12.0	12,854	16.3



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 5/5/04

Summary of Tre	nds in C	Growth ar	d Anen	nia Ind	icators b	y Race	Ethnicity	(3)	
Race/Ethnicity and Year			Heigh	nt and W	eight				
			Short Stature (4)	Under- weight (5)			Anemia Low Hb/Hct — (6)		
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	Number	%	
Total All Race/Ethnic Groups	2003	490,014	6.4	2.7	18.4	21.7	437,622	12.7	
	2002	487,486	6.7	2.7	18.1	20.8	429,702	12.0	
	2001	454,349	6.8	2.7	17.8	19.9	397,255	12.4	
	2000	555,211	7.0	2.8	17.7	19.7	473,293	12.6	
	1999	579,775	6.8	2.8	17.6	18.9	498,396	11.9	
	1998	527,057	6.9	2.8	17.2	17.6	458,165	12.1	
	1997	489,939	7.0	3.2	16.5	16.4	429,207	12.9	
	1996	418,996	7.5	3.2	16.3	15.7	401,444	13.1	
	1995	389,137	8.1	3.3	16.2	15.1	371,725	13.7	
	1994	355,146	8.6	3.4	15.8	14.4	344,767	13.8	



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.